

INTERPRETATIONS OF OCTONION WAVE EQUATIONS

P. S. Bisht⁽¹⁾ , Bimal Pandey⁽¹⁾ and O. P. S. Negi⁽²⁾ *

February 1, 2008

⁽¹⁾Department of Physics,
Kumaun University,
S. S. J. Campus,
Almora - 263601 (Uttarakhand) India

⁽²⁾Institute of Theoretical Physics,
Chinese Academy of Sciences
KITP Building Room No.- 6304
Hai Dian Qu Zhong Guan Chun Dong Lu
55 Hao , Beijing - 100080, P.R.China

Email - ps_bisht123@rediffmail.com,
ops_negi@yahoo.co.in

Abstract

The interpretations of octonion wave equations in eight dimensional space-time have been discussed. We have made an attempt to discuss the octonion field equation as the equation of motion for particles carrying simultaneously electric and magnetic charges (i.e. dyons) in external and internal spaces respectively. It has been concluded that the component of octonion potential wave equations behaves neither the generalized electromagnetic fields of monopoles nor the dyons. Rather, they have the mixed behaviour of electromagnetic fields associated with the electric and magnetic charges in external and internal spaces. We have also made an attempt to investigate the split octonion wave equation and its interpretation in classical electrodynamics and accordingly the consistent and compact forms of eight-dimensional potential and current equation of dyons are obtained in terms of Zorn's vector matrix realization of split octonions. Visualizing the external four space as the localization space for tachyons, it is shown that the split octonion wave equation reduces to the Maxwell's equation (field equation) for bradyons in R^4 - space as well as that for tachyons in T^4 - space in the absence of other.

*Permanent Address- Department of Physics, Kumaun University, S. S. J. Campus, Almora -263601 (U.A.) INDIA

1 INTRODUCTION

Octonions were first introduced in Physics by Jordon, Von Neuman and Wigner[1], who investigated a new finite Hilbert space, on replacing the complex numbers by Octonions[2]. According to celebrated Hurwitz theorem [3] there exists four-division algebra consisting of \mathbb{R} (real numbers), \mathbb{C} (complex numbers), \mathbb{H} (quaternions) and \mathcal{O} (octonions). All four algebras are alternative with antisymmetric associators. In 1961 Pais [4] pointed out a striking similarity between the algebra of interactions and the split octonion algebra. Recently, some attention has been directed to octonions in theoretical physics with the hope of extending the 3+1 space-time framework of the theory to eight dimensions to accommodate the ever increasing quantum numbers and internal symmetries assigned to elementary particles and gauge fields. A lot of literature is available [5, 6, 7, 8, 9, 10, 11] on the applications of octonions to interpret wave equation, Dirac equation, and the extension of octonion non-associativity to physical theories. It is assumed there that octonion wave equation contains four-dimensional internal space of mass parameters while the external four dimensional space has been interpreted as the four generators of fermions within the quark-lepton symmetry. The ingenious work, in this direction is done by Günaydin and Gürsey [12, 13], to formulate quark models and colour gauge theory in terms of split octonions. The $SU(3)$ group appears as the automorphism group of octonion representation leaving the complex subspace and the scalar product invariant. This approach is extended by many [14, 15, 16] to investigate the role of octonions and division algebra in unified gauge theories, higher dimensional theories of supersymmetry and super strings. Octonions are also used by Buoncristiani [17] in writing Yang-Mill's field equation in a simpler form. The extension of quaternion matrices to octonions for their interpretation in non-Riemannian geometry is described by Marques et al in a series of papers [18, 19, 20, 21, 22].

During the past few years, there has been considerable interest in higher-dimensional kinematical models [23, 24] for a proper and unified representation of relativistic object, bradyonic as well as tachyonic (including those with internal structure). It has been speculated [25, 26, 27] that the problem of representation and localization of tachyonic objects may be solved only with the extension of four-dimensional Minkowski space to higher-dimensional space-time. It has already been discussed in a series of papers [28, 29, 30, 31, 32, 33] that the true localization space for the representation of tachyons is T^4 -space with one space and three time coordinates while that for bradyons is the usual R^4 -space with one time and three space coordinates. The unified eight dimensional space time has been discussed earlier [32, 33] as the unified space of bradyons and tachyons i.e. $R^8 = R^4 \cup T^4$. The two R^4 - and T^4 -localization spaces are considered as external (internal) space for bradyons (tachyons) and vice versa. Moreover, the built-in duality associated with the combination of symmetries of R^4 - and T^4 -subspaces is useful to understand the problem of quark confinement in quantum chromodynamics where the role of tricolors would be played by three time coordinates for bradyons and those of three space coordinates for tachyons.

In order to interpret octonion wave equations in eight dimensional space- time, in the present paper, we have made an attempt to discuss the octonion field equation as the equation of motion for particles carrying simultaneously electric and magnetic (monopole [34]) charges (i.e. dyons [35]). Starting with the regularity condition in the octonion field equation we have developed consistent and compact formulation of eight-dimensional potential and current equations of dyons. It has been demonstrated that the component of octonion potential wave equations behaves neither the generalized electromagnetic fields of monopoles nor the dyons. Rather, they have the mixed behaviour of electromagnetic fields associated with the electric and magnetic charges in external and internal

spaces.

After decomposing the octonion wave equation in to two quaternion-valued wave equations, it is shown that the two different spaces demonstrate the separate wave equations for electromagnetic fields. Visualizing the external four spaces as the localization space for bradyons and the internal space as the localization space for tachyons, it is shown that the octonion wave equation reduces to the Maxwell's equation (field equation) for bradyons in R^4 - space as well as that for tachyons in T^4 - space. It has also been emphasized that the quaternionic decomposition of octonion wave equation gives rise to the equations of dyons and tachyonic dyons, when we take them as combination of bi-quaternion instead of real quaternion in external and internal four-dimensional spaces of eight dimensional space time.

Then we have made an attempt to investigate the split octonion wave equation and its interpretation in classical electrodynamics. Split octonion electrodynamics has been deduced in terms of Zorn's vector matrix realization by describing electrodynamics potential, current and other dynamical quantities as octonion variables. Also the consistent and compact forms of eight-dimensional potential and current equation of dyons are obtained in terms of Zorn's vector matrix realization of split octonions. It has been shown that the octonion valued potential wave equation behaves neither the generalized electromagnetic fields of monopoles nor of the dyons. Rather, they have the mixed behaviour of electromagnetic fields associated with the electric and magnetic charges in external and internal spaces. At last, it is shown that split octonion Zorn's vector realization reproduces two different spaces to demonstrate the separate wave equations for electromagnetic fields. Visualizing the external four space as the localization space for tachyons, it is shown that the octonion wave equation, when expressed in terms of split octonions, reduces to the Maxwell's equation (field equation) for bradyons in R^4 - space as well as that for tachyons in T^4 - space in the absence of other.

2 OCTONION DEFINITION

An octonion x is expressed as a set of eight real numbers

$$x = e_0x_0 + e_1x_1 + e_2x_2 + e_3x_3 + e_4x_4 + e_5x_5 + e_6x_6 + e_7x_7 = e_0x_0 + \sum_{A=1}^7 e_Ax_A \quad (1)$$

where $e_A (A = 1, 2, \dots, 7)$ are imaginary octonion units and e_0 is the multiplicative unit element. Set of octets $(e_0, e_1, e_2, e_3, e_4, e_5, e_6, e_7)$ are known as the octonion basis elements and satisfy the following multiplication rules

$$e_0 = 1; \quad e_0e_A = e_Ae_0 = e_A; \quad e_Ae_B = -\delta_{AB}e_0 + f_{ABC}e_C. \quad (A, B, C = 1, 2, \dots, 7) \quad (2)$$

The structure constants f_{ABC} is completely antisymmetric and takes the value 1 for following combinations,

$$f_{ABC} = +1; \quad \forall(ABC) = (123), (471) (257) (165) (624) (543) (736). \quad (3)$$

It is to be noted that the summation convention is used for repeated indices. Here the octonion algebra \mathcal{O} is described over the algebra of rational numbers having the vector space of dimension 8. We now

get the following relations among octonion basis elements

$$\begin{aligned} [e_A, e_B] &= 2 f_{ABC} e_C; \\ \{e_A, e_B\} &= -2 \delta_{AB} e_0; \\ e_A(e_B e_C) &\neq (e_A e_B) e_C \end{aligned} \quad (4)$$

where brackets $[]$ and $\{ \}$ are used respectively for commutation and the anti commutation relations while δ_{AB} is the usual Kronecker delta-Dirac symbol. Octonion conjugate is defined as

$$\overline{x} = e_0 x_0 - e_1 x_1 - e_2 x_2 - e_3 x_3 - e_4 x_4 - e_5 x_5 - e_6 x_6 - e_7 x_7 = e_0 x_0 - \sum_{A=1}^7 e_A x_A \quad (5)$$

where we have used the conjugates of basis elements as $\overline{e_0} = e_0$ and $\overline{e_A} = -e_A$. Hence an octonion can be decomposed in terms of its scalar ($Sc(x)$) and vector ($Vec(x)$) parts as

$$Sc(x) = \frac{1}{2}(x + \overline{x}); \quad Vec(x) = \frac{1}{2}(x - \overline{x}) = \sum_{A=1}^7 e_A x_A. \quad (6)$$

Conjugates of product of two octonions and its own are described as

$$\overline{(xy)} = \overline{y} \overline{x}; \quad \overline{(\overline{x})} = x. \quad (7)$$

while the scalar product of two octonions is defined as

$$\langle x, y \rangle = \frac{1}{2}(x \overline{y} + y \overline{x}) = \frac{1}{2}(\overline{x} y + \overline{y} x) = \sum_{\alpha=0}^7 x_\alpha y_\alpha. \quad (8)$$

The norm $N(x)$ and inverse x^{-1} (for a nonzero x) of an octonion are respectively defined as

$$\begin{aligned} N(x) = x \overline{x} = \overline{x} x &= \sum_{\alpha=0}^7 x_\alpha^2 e_0; \\ x^{-1} &= \frac{\overline{x}}{N(x)} \implies x x^{-1} = x^{-1} x = 1. \end{aligned} \quad (9)$$

The norm $N(x)$ of an octonion x is zero if $x = 0$, and is always positive otherwise. It also satisfies the following property of normed algebra

$$N(xy) = N(x)N(y) = N(y)N(x). \quad (10)$$

Equation (4) shows that octonions are not associative in nature and thus do not form the group in their usual form. Non-associativity of octonion algebra \mathcal{O} is provided by the associator $(x, y, z) =$

$(xy)z - x(yz) \forall x, y, z \in \mathcal{O}$ defined for any 3 octonions. If the associator is totally antisymmetric for exchanges of any 2 variables, i.e. $(x, y, z) = -(z, y, x) = -(y, x, z) = -(x, z, y)$, then the algebra is called alternative. Hence, the octonion algebra is neither commutative nor associative but, is alternative.

3 OCTONION WAVE EQUATION

In order to write the Octonion wave equation, let us define the differential octonion D as

$$D = \sum_{\mu=0}^7 e_{\mu} D_{\mu} \quad (11)$$

where D_{μ} are described as the components of differential operator in eight dimensional representation. Here we assume the eight - dimensional space as the combination of two (external and internal) four-dimensional spaces. As such, we describe a function of octonion variable as

$$\mathcal{F}(X) = \sum_{\mu=0}^7 e_{\mu} f_{\mu}(X) = f_0 + e_1 f_1 + e_2 f_2 + \dots + e_7 f_7 \quad (12)$$

where f_{μ} are scalar functions. Since octonions are neither commutative nor associative, one has to be very careful to multiply the octonion either from left or from right in terms of regularity conditions.

As such, a function $\mathcal{F}(X)$ of an octonion variable $X = \sum_{\mu=0}^7 e_{\mu} X_{\mu}$ is left regular at X if and only if $\mathcal{F}(X)$ satisfies the condition

$$D\mathcal{F}(X) = 0. \quad (13)$$

Similarly, a function $G(X)$ is a right regular if and only if

$$G(X)D = 0 \quad (14)$$

where $G(X) = g_0 + g_1 e_1 + g_2 e_2 + \dots + g_7 e_7$. Then we get

$$D\mathcal{F} = I_0 + I_1 e_1 + I_2 e_2 + I_3 e_3 + I_4 e_4 + I_5 e_5 + I_6 e_6 + I_7 e_7 \quad (15)$$

where

$$\begin{aligned}
I_0 &= \partial_0 f_0 - \partial_1 f_1 - \partial_2 f_2 - \partial_3 f_3 - \partial_4 f_4 - \partial_5 f_5 - \partial_6 f_6 - \partial_7 f_7; \\
I_1 &= \partial_0 f_1 + \partial_1 f_0 + \partial_2 f_3 - \partial_3 f_2 + \partial_6 f_5 - \partial_5 f_6 - \partial_7 f_4 + \partial_4 f_7; \\
I_2 &= \partial_0 f_2 + \partial_2 f_0 + \partial_3 f_1 - \partial_1 f_3 + \partial_4 f_6 - \partial_6 f_4 - \partial_7 f_5 + \partial_5 f_7; \\
I_3 &= \partial_0 f_3 + \partial_3 f_0 + \partial_1 f_2 - \partial_2 f_1 + \partial_6 f_7 - \partial_7 f_6 + \partial_5 f_4 - \partial_4 f_5; \\
I_4 &= \partial_0 f_4 + \partial_4 f_0 + \partial_3 f_5 - \partial_5 f_3 - \partial_2 f_6 + \partial_6 f_2 - \partial_1 f_7 + \partial_7 f_1; \\
I_5 &= \partial_0 f_5 + \partial_5 f_0 + \partial_1 f_6 - \partial_6 f_1 + \partial_7 f_2 - \partial_2 f_7 - \partial_3 f_4 + \partial_4 f_3; \\
I_6 &= \partial_0 f_6 + \partial_6 f_0 - \partial_1 f_5 + \partial_5 f_1 + \partial_2 f_4 - \partial_4 f_2 - \partial_3 f_7 + \partial_7 f_3; \\
I_7 &= \partial_0 f_7 + \partial_7 f_0 + \partial_1 f_4 - \partial_4 f_1 + \partial_2 f_5 - \partial_5 f_2 - \partial_6 f_3 + \partial_3 f_6.
\end{aligned} \tag{16}$$

The regularity condition (13) may now be considered as homogeneous octonion wave equation for octonion variables without sources. On the other hand, equation (15) is considered as the inhomogeneous wave equation

$$D\mathcal{F} = I \tag{17}$$

where I is again an octonion. Similarly, we may also write the homogeneous as well as inhomogeneous octonion wave equations on using the right regularity condition (14). We may now interpret these octonion wave equations as the classical wave (field) equations of physical variables.

3.1 POTENTIAL EQUATION FOR ELECTROMAGNETIC FIELDS WITH SOURCES

Let us consider the case of generalized electromagnetic fields of dyons (particles carrying simultaneous existence of electric and magnetic charges). We may now define an octonion valued potential, in eight dimensional formalism as the combinations of two four-dimensional spaces, as follows

$$\emptyset = \sum_{\mu=0}^7 e_\mu \emptyset_\mu = \sum_{\mu=0}^3 e_\mu A_\mu + \sum_{\nu=4}^7 e_\nu B_\nu; \tag{18}$$

where \emptyset is the octonion potential and A_μ and B_ν are assumed respectively as the electric and magnetic four - potentials associated with the electric and magnetic charges of dyons [36]. As such, we may write the wave equation for octonion potential variable for simultaneous existence of electric and magnetic charges on a particle (namely dyons) in the following manner ,

$$\overline{D}\emptyset = F \tag{19}$$

where

$$\overline{D} = F \sum_{\mu=0}^7 \overline{e}_\mu D_\mu = e_0 D_0 - \sum_{A=1}^7 e_A D_A; \quad F = \sum_{\mu=0}^7 F_\mu e_\mu. \tag{20}$$

The coefficients of octonion F are then given by

$$\begin{aligned}
F_0 &= \partial_0\emptyset_0 + \partial_1\emptyset_1 + \partial_2\emptyset_2 + \partial_3\emptyset_3 + \partial_4\emptyset_4 + \partial_5\emptyset_5 + \partial_6\emptyset_6 + \partial_7\emptyset_7; \\
F_1 &= \partial_0\emptyset_1 - \partial_1\emptyset_0 + \partial_3\emptyset_2 - \partial_2\emptyset_3 + \partial_7\emptyset_4 - \partial_4\emptyset_7 + \partial_5\emptyset_6 - \partial_6\emptyset_5; \\
F_2 &= \partial_0\emptyset_2 - \partial_2\emptyset_0 + \partial_1\emptyset_3 - \partial_3\emptyset_1 + \partial_6\emptyset_4 - \partial_4\emptyset_6 + \partial_7\emptyset_5 - \partial_5\emptyset_7; \\
F_3 &= \partial_0\emptyset_3 - \partial_3\emptyset_0 + \partial_2\emptyset_1 - \partial_1\emptyset_2 + \partial_4\emptyset_5 - \partial_5\emptyset_4 + \partial_7\emptyset_6 - \partial_6\emptyset_7; \\
F_4 &= \partial_0\emptyset_4 - \partial_4\emptyset_0 - \partial_3\emptyset_5 + \partial_5\emptyset_3 + \partial_2\emptyset_6 - \partial_6\emptyset_2 + \partial_1\emptyset_7 - \partial_7\emptyset_1; \\
F_5 &= \partial_0\emptyset_5 - \partial_5\emptyset_0 - \partial_1\emptyset_6 + \partial_6\emptyset_1 - \partial_7\emptyset_2 + \partial_2\emptyset_7 + \partial_3\emptyset_4 - \partial_4\emptyset_3; \\
F_6 &= \partial_0\emptyset_6 - \partial_6\emptyset_0 + \partial_1\emptyset_5 - \partial_5\emptyset_1 - \partial_2\emptyset_4 + \partial_4\emptyset_2 + \partial_3\emptyset_7 - \partial_7\emptyset_3; \\
F_7 &= \partial_0\emptyset_7 - \partial_7\emptyset_0 - \partial_1\emptyset_4 + \partial_4\emptyset_1 - \partial_2\emptyset_5 + \partial_5\emptyset_2 + \partial_6\emptyset_3 - \partial_3\emptyset_6.
\end{aligned} \tag{21}$$

We get $F_0 = 0$ in the Euclidean space $(+++++)$ due to eight dimensional Lorentz gauge condition (i.e the combination of two gauges associated with electric and magnetic potentials). Thus, one dimensional octonion representation is identical to eight dimensional spaces over the field of real numbers. It is isomorphic to four-dimensional space representation over the field of complex variables which is equivalent to two-dimensional space representation over quaternion field variables. Similarly, one dimensional quaternion space is isomorphic to four-dimensional space over the field of real numbers which is identical to two-dimensional space over the field of complex numbers. So, an octonionic potential may also be described as the combination of two quaternion potentials in the following manner;

$$\emptyset = \emptyset_a + e_7\emptyset_b; \quad D = D_a + e_7D_b; \quad F = F_a + e_7F_b \tag{22}$$

where $\emptyset_a, \emptyset_b, D_a, D_b, F_a$ and F_b are quaternion variables described as

$$\begin{aligned}
\emptyset_a &= \emptyset_0 + e_1\emptyset_1 + e_2\emptyset_2 + e_3\emptyset_3; \\
\emptyset_b &= \emptyset_7 + e_1\emptyset_4 + e_2\emptyset_5 + e_3\emptyset_6; \\
D_a &= \partial_0 + e_1\partial_1 + e_2\partial_2 + e_3\partial_3; \\
D_b &= \partial_7 + e_1\partial_4 + e_2\partial_5 + e_3\partial_6; \\
F_a &= F_0 + e_1F_1 + e_2F_2 + e_3F_3; \\
F_b &= F_7 + e_1F_4 + e_2F_5 + e_3F_6.
\end{aligned} \tag{23}$$

Here e_1, e_2 , and e_3 ; three quaternion units; satisfy the multiplication rule $e_j e_k = -\delta_{jk} + \varepsilon_{jkl} e_l \quad \forall (j, k, l = 1, 2, 3)$. If we replace the octonion unit e_7 by imaginary unit $i = \sqrt{-1}$, commuting with all the octonion basis elements e_A , the resultant theory of octonions become the theory of bi-quaternion variables where generalized fields of dyons are written in compact, covariant and consistent way [37]. We may now consider the following mapping for different four-dimensional spaces,

$$\begin{aligned}
\partial_7 &\rightarrow \partial'_0, & \emptyset_7 &\rightarrow B_0, & \emptyset_0 &\rightarrow A_0; \\
\partial_4 &\rightarrow \partial'_1, & \emptyset_4 &\rightarrow B_1, & \emptyset_1 &\rightarrow A_1; \\
\partial_5 &\rightarrow \partial'_2, & \emptyset_5 &\rightarrow B_2, & \emptyset_2 &\rightarrow A_2; \\
\partial_6 &\rightarrow \partial'_3, & \emptyset_6 &\rightarrow B_3, & \emptyset_3 &\rightarrow A_3.
\end{aligned} \tag{24}$$

Then we get

$$\begin{aligned}
F_j &= \partial_0 A_j - \partial_j A_0 + (\vec{\nabla} \times \vec{A})_j + \partial'_0 B_j - \partial'_j B_0 + (\vec{\nabla} \times \vec{B})_j; \\
F_{j+3} &= \partial_0 B_j - \partial_j B_0 + (\vec{\nabla} \times \vec{B})_j + \partial'_0 A_j - \partial'_j A_0 + (\vec{\nabla}' \times \vec{A})_j; \\
F_7 &= \partial_0 B_0 - \partial_1 B_1 - \partial_2 B_2 - \partial_3 B_3 - (\partial'_0 A_0 + \partial'_1 A_1 + \partial'_2 A_2 + \partial'_3 A_3).
\end{aligned} \tag{25}$$

Hence equation (25) shows that F_0 , F_j , F_{j+3} and F_7 correspond to the components of electric and magnetic fields obtained in terms of components of two four-potentials (A_μ and B_μ) in internal and external spaces. If these two spaces are completely disjoint spaces, we get only F_j ; and F_0 vanishing due to Lorentz gauge condition; while F_{j+3} and F_7 do not occur. Here prime derivatives are associated with the second four-dimensional structure of eight potential. In equation (25), we find that F_j is made up of the j^{th} components of electric and magnetic fields. Here the first term $\partial_0 A_j - \partial_j A_0$ describes the electric field and second term $(\vec{\nabla} \times \vec{A})_j$ corresponds to the magnetic field in four-dimensional (we call it as the external) space. In the light of the duality mechanism between electric and magnetic fields (and for electric $\{A_\mu\}$ and magnetic $\{B_\mu\}$ four-potentials), the third term $\partial_0 B_j - \partial_j B_0$ of F_j is equivalent to the magnetic field while the last term $(\vec{\nabla} \times \vec{B})_j$ denotes the electric field in other four-dimensional (let us call it as internal or magnetic) space. On mixing of these two spaces, $\partial_0 A_j - \partial_j A_0 + (\vec{\nabla} \times \vec{A})_j$ resembles with the expression of generalized electric field and $\partial'_0 B_j - \partial'_j B_0 + (\vec{\nabla}' \times \vec{B})_j$ as that of the generalized magnetic fields of dyons [36, 37]. So, it looks awkward to combine external (internal) space of electric charge (magnetic monopole) to the internal (external) space of magnetic monopole (electric charge) to interpret F_j . As such, we may conclude that the equation (25) does not represent the true generalization of potential equation for the generalized electromagnetic fields of dyons. Rather, we have obtained the mixed behaviour of electric and magnetic charges in internal and external spaces. So, octonion wave equation of electromagnetic potential faces difficulties and hence needs modification to interpret it consistently.

3.2 CURRENT EQUATION

We now Analyse the octonion wave equation (17) as the current equation in eight dimensional representations i.e.

$$D \quad F = \quad S \tag{26}$$

where

$$\begin{aligned}
S_0 &= \partial_0 F_0 - \partial_1 F_1 - \partial_2 F_2 - \partial_3 F_3 - (\partial'_1 F_4 + \partial'_2 F_5 + \partial'_3 F_6 + \partial'_0 F_7); \\
S_j &= \partial_0 F_j + \partial_j F_0 + (\vec{\nabla} \times \vec{F})_j - \partial'_0 F_{j+3} + \partial'_j F_7 - (\vec{\nabla}' \times \vec{F})_{j+3}; \\
S_{j+3} &= \partial'_0 F_j + \partial'_j F_0 - (\vec{\nabla}' \times \vec{F})_j - \partial_0 F_{j+3} - \partial_j F_7 - (\vec{\nabla} \times \vec{F})_{j+3}; \\
S_7 &= \partial_1 F_4 + \partial_2 F_5 + \partial_3 F_6 + \partial_0 F_7 + (\partial'_0 F_0 - \partial'_1 F_1 - \partial'_2 F_2 - \partial'_3 F_3).
\end{aligned} \tag{27}$$

Thus, the current equation for octonion variables describes [38] the generalized structure of differential equations (Maxwell's equations) in eight - dimensional space-time. Four- dimensional reduction of these differential equations may be visualized as the Maxwell's equations in internal and external spaces. So, it is difficult to explain the various terms associated with the eight parameters S_Λ ($\Lambda = 0, 1, 2, 3, \dots, 7$) correctly. Silagade [38] provided the interpretation of the homogeneous octonion wave equation $DF = 0$ as equivalent to one of the pair of seven-dimensional Maxwell's equations and the second pair of seven dimensional Maxwell's equations may be obtained on applying the duality transformations between electric and magnetic fields therein. Similarly, Gamba and Gogberashvili [8] also tried to explain the structure of eight parameters S_Λ ($\Lambda = 0, 1, 2, 3, \dots, 7$) but the consistent justification needs modifications. Hence, octonion wave equation (26) may be identified rather a current equation or octonion field equation in eight dimensional spaces for mixed structural behaviour of sources (the electric and magnetic) instead of dyons. In other words we may say that the equation (26) can not be described as the true field equation of dyons. Here S_0 and S_7 may be taken vanishing due to equations of continuity in internal and external spaces.

3.3 QUATERNION DECOMPOSITION

We may decompose an octonion in terms of two-quaternions. So, let us decompose octonion wave equation for potential in terms of two quaternions as

$$\overline{D}\emptyset = (\overline{D}_a - e_7 D_b)(\emptyset_a + e_7 \emptyset_b) = (\phi + e_7 \varphi) = F. \tag{28}$$

Here

$$\phi = \overline{D}_a \emptyset_a + \emptyset_b \tilde{D}_b = (D_0 - D_1 e_1 - D_2 e_2 - D_3 e_3) \emptyset_a + \emptyset_b (\overleftarrow{D}_7 - \overleftarrow{D}_4 e_1 - \overleftarrow{D}_5 e_2 - \overleftarrow{D}_6 e_3) \tag{29}$$

where $[\overleftarrow{D}_r, (r = 4, 5, 6, 7)]$ represents the partial differential of \emptyset_b from right to left, \tilde{D}_b denotes the quaternion conjugation ($\tilde{e}_0 = e_0$; $\tilde{e}_j = -e_j \quad \forall j = 1, 2, 3$) from right to left i.e.

$$\overline{D}_a = D_0 - D_1 e_1 - D_2 e_2 - D_3 e_3;$$

$$\tilde{\overline{D}}_a = D_0 + D_1 e_1 + D_2 e_2 + D_3 e_3 = D_1;$$

$$\tilde{D}_b = D_7 - D_4 e_1 - D_5 e_2 - D_6 e_3$$

and

$$\varphi = \overline{\tilde{D}_a} \quad \emptyset_b - \emptyset_a D_b = D_a \emptyset_b - \emptyset_a D_b. \quad (30)$$

Here an octonion-valued potential is described as the sum of two quaternions by Cayley-Dickson doubling process. Thus, we observe that the octonion potential wave equation rules out the existence of isolated of magnetic monopole and provides a mixed structural behaviour of electromagnetic fields containing both charges simultaneously. In other words, we may say that it is a generalized field equation of electromagnetic fields associated with a particle carrying simultaneous existence of electric and magnetic charges (namely dyons). The generalized electromagnetic fields of dyons are symmetric and dual invariant in terms of two four potentials. So, the possibility of monopoles (or dyons) in non Abelian or supersymmetric gauge theories is directly linked with the existence of second four potential which is hidden in external space and present in internal space while the electric four potential is the consequence of our external space and seems to be hidden in internal space.

Similarly we may express the current equation given by equation (26) in terms of two quaternions as

$$DF = (D_a + e_7 D_b)(g + e_7 h) = S = s_a + e_7 s_b \quad (31)$$

where

$$\begin{aligned} s_a &= D_a g - h D_b = D_a \overline{D}_b \emptyset_a + \emptyset_b D_b \overline{D}_b \\ s_b &= \tilde{D}_a h + g D_b = \overline{D}_a D_a \emptyset_b + \emptyset_b \overline{D}_b D_b. \end{aligned} \quad (32)$$

Thus $S = s_a + e_7 s_b$ is the octonion expression for generalized current density. Similarly, we may express the generalized force and generalized field tensor density in terms of two quaternions.

3.4 LOCALIZATION SPACES OF BRADYONS AND TACHYONS

We have described the octonion (eight dimensional) space is made of two quaternion (namely the external and internal four dimensional) spaces. Let us suppose the external four space as the usual Minkowski (or Euclidean) $R^4 \Rightarrow M(1, 3)$ - space (consisting one time and three space coordinates). This space has been named [23, 24] as. the localization space of bradyons (particles traveling slower than light \rightarrow subluminal particles). Accordingly, we describe the internal four dimensional space as the $T^4 \Rightarrow M(3, 1)$ - space (consisting three time and one space coordinates). The possibility of such space is explored [23, 24] as the localization space of tachyons (particles traveling faster than light \rightarrow superluminal particles). Hence an octonion eight dimensional space is described as the unified

space containing both external $R^4 \Rightarrow M(1, 3)$ and internal $T^4 \Rightarrow M(3, 1)$ four - dimensional spaces. In other words we may identify the octonion space as the unified localization space for the description of bradyons and tachyons. It is because that the bradyons are the objects of our four dimensional world for which the Cauchy's data are described on the plane $\{t = 0\}$ while tachyons are localized in time and their Cauchy's data lies on the plane $\{x = 0\}$. Hence, the eight dimensional space of octonion variables described above is the unified localization space for bradyons and tachyons and hence be expressed as $R^8 = R^4 \cup T^4$. The T^4 -space is characterized as the hidden space for bradyons while R^4 -space is identified as the hidden space for tachyons. We may also interpret that the R^4 - space is visualized as the internal space for tachyons (and external space for bradyons) and correspondingly the T^4 - space is the internal space for bradyons (and external space for tachyons). First quaternion variable of equation (31) maps to four-dimensional space for bradyons i.e. $R^4 \simeq (t, \vec{r})$ while the second quaternion variable gives rise to four-dimensional space for tachyons i.e. $T^4 \simeq (r, \vec{t})$. In the absence of second quaternion, equation (32) reduces to

$$s_a = D_a \bar{D}_a \emptyset_a \Rightarrow \left(\frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} - \frac{\partial^2}{\partial z^2} \right) \phi = \square(R) \phi \quad (33)$$

which resembles to Cauchy-Feuter equation of quaternion variables in $R^4 \simeq (t, \vec{r})$ space and thus describes the usual Maxwell's equations in electromagnetic fields in R^4 - space. On the other hand in absence of first quaternion, equation (32) reduces to

$$s_b = \emptyset_b \quad \bar{D}_b D_b \Rightarrow \varphi \left(\frac{\partial^2}{\partial r^2} - \frac{\partial^2}{\partial t_x^2} - \frac{\partial^2}{\partial t_y^2} - \frac{\partial^2}{\partial t_z^2} \right) = \varphi \overleftarrow{\square}(T) \quad (34)$$

which also describes the Cauchy - Feuter equation of quaternion variables in $T^4 \simeq (r, \vec{t})$ space and coincides with the Maxwell's equations of superluminal photons in T^4 - space. Equation (33) is the left regular while equation (34) is the right regular. hence, the regularity conditions are changes for internal and external spaces. In case of bi-quaternion, equations (33) and (34) characterize respectively to the equations of dyons for subluminal and superluminal electromagnetic fields. In general (i.e. octonionic space) describes the unified structure of subluminal and superluminal objects. As such, the octonionic eight dimensional representation is the unified picture of bradyons and tachyons and the octonionic current equations thus reproduce two different kinds of Maxwell's equations in external and internal spaces in order to explain the simultaneous description of bradyons and tachyons. Here, we have developed the octonion wave equation and theory of octonion field variables in compact and simpler manner. So, in order to overcome the non-associativity, it is necessary to decompose them in terms of two quaternions isomorphic to two different four-dimensional spaces. The compact and simpler form of octonion field equation provide an unified model of the theories of subluminal and superluminal electromagnetic fields in view of the localizability of bradyons and tachyons in consistent manner.

4 SPLIT OCTONION WAVE EQUATION

The split octonions are a non associative extension of quaternions (or the split quaternions). They differ from the octonion in the signature of quadratic form. The split octonions have a signature (4, 4) whereas the octonions have positive signature (8, 0). The Cayley algebra of octonions over the field of complex numbers $\mathbb{C}_C = \mathbb{C} \otimes \mathbb{C}$ is visualized as the algebra of split octonions with basis elements

$u_0 = \frac{1}{2}(e_0 + ie_7)$; $u_0^* = \frac{1}{2}(e_0 - ie_7)$, $u_j = \frac{1}{2}(e_j + ie_{j+3})$, $u_j^* = \frac{1}{2}(e_j - ie_{j+3})$ ($j = 1, 2, 3; i = \sqrt{-1}$) as the bi-valued (or bi - dimensional) representations of quaternion units e_0, e_1, e_2, e_3 . The split Cayley (octonion) algebra is thus expressed in terms of 2×2 Zorn's vector matrices components of which are scalar and vector parts of a quaternion. As such, we may also write an arbitrary split octonion A in terms of following 2×2 Zorn's vector matrix realizations [39] as

$$A = au_0^* + bu_0 + x_i u_i^* + y_i u_i = \begin{pmatrix} a & -\vec{x} \\ \vec{y} & b \end{pmatrix}. \quad (35)$$

Split octonion conjugation of equation (35) is then described as,

$$\overline{A} = au_0 + bu_0^* - x_i u_i^* - y_i u_i = \begin{pmatrix} b & \vec{x} \\ -\vec{y} & a \end{pmatrix}. \quad (36)$$

The norm of A is defined as,

$$\overline{A}A = A\overline{A} = (ab + \vec{x} \cdot \vec{y})\hat{1} \quad (37)$$

where $\hat{1}$ is the identity element given by $\hat{1} = 1u_0 + 1u_0^*$. Any four - vector A_μ (complex or real) can equivalently be written in the following Zorn's matrix realization as

$$Z(A) = \begin{pmatrix} x_4 & -\vec{x} \\ \vec{y} & y_4 \end{pmatrix} \quad (38)$$

and

$$Z(\overline{A}) = \begin{pmatrix} x_4 & \vec{x} \\ -\vec{y} & y_4 \end{pmatrix}. \quad (39)$$

In equation (38, 39), putting $\vec{x} = \vec{y}$, we get the equivalent matrix realization for a four vector in bi-valued four-dimensional Euclidean space - time. In order to develop the the eight dimensional split octonion kinematics, we start with the following definition of the split octonion differential operator D and its conjugate \overline{D} in terms of 2×2 Zorn's matrix realizations,

$$D = \begin{pmatrix} \partial_4 & -\vec{\nabla} \\ \vec{\nabla} & \partial'_4 \end{pmatrix} \quad (40)$$

$$\overline{D} = \begin{pmatrix} \partial_4 & \vec{\nabla} \\ -\vec{\nabla} & \partial'_4 \end{pmatrix} \quad (41)$$

with the assumption that the primed variables are represented in internal space whereas the unprimed variables are defined in external four - dimensional space.

4.1 POTENTIAL EQUATION

Let us express an octonion potential in the following form

$$\emptyset = A_4 u_0 * + B_4 u_0 + B_i u_i * + A_i u_i \quad (42)$$

where $\{A_\mu\} = (A_4, A_j)$ and $\{B_\mu\} = (B_4, B_j)$ are two four - potentials respectively associated with electric and magnetic charges. Equation (42) may then be expressed in terms of split octonion 2×2 Zorn's vector matrix realization as

$$\emptyset = \begin{pmatrix} A_4 & -\vec{B} \\ \vec{A} & B_4 \end{pmatrix} \quad (43)$$

where $A_4 = i \emptyset_e$ (e denotes the electric charge) and $B_4 = i \emptyset_g$ (g denotes the magnetic charge). Here \vec{A} and A_4 are assumed to be the components of electric four - potential $A_\mu = (\vec{A}, i \emptyset_e)$. Similarly \vec{B} and B_4 are considered as the components of magnetic four-potential $B_\mu = (\vec{B}, i \emptyset_g)$. We have designed the eight dimensional space spanned by split octonion basis elements in terms of two four - dimensional space (namely the external and internal space). Here also the electric four-potential is described in external four-dimensional space while the magnetic four - potential has been considered in internal four dimensional spaces. Since electric charge and magnetic monopoles are dual to each other ,we may interpret these two spaces as dual to each other. The internal space of electric charge may be identified as the external space of monopole while the external space of electric charge will be visualized as the internal space of magnetic monopole or vice versa. Accordingly, the electric charge is considered in external space and magnetic charge in internal space. Hence, eight dimensional spaces have the built in duality where a bradyonic monopole plays the role of tachyonic electric charge while the tachyonic monopole looks like a bradyonic electric charge or vice versa. Eight dimensional spaces have been considered as the unification of R^4 - (bradyonic) and T^4 - (tachyonic) spaces.

Let us now write the the inhomogeneous octonion wave equation in its split form as

$$\overline{D}\emptyset = \begin{pmatrix} \partial_4 A_4 + \vec{\nabla} \cdot \vec{A} & -(\partial_4 \vec{B} - \vec{\nabla} B_4 - \vec{\nabla} \times \vec{A}) \\ \partial'_4 \vec{A} - \vec{\nabla}' A_4 - \vec{\nabla}' \times \vec{B} & \partial'_4 B_4 + \vec{\nabla}' \cdot \vec{B} \end{pmatrix} = F \quad (44)$$

where F is a field like electromagnetic field in split octonion 2×2 Zorn's vector matrix realization as

$$F = \begin{pmatrix} f_4 & -\vec{f} \\ \vec{f}' & f'_4 \end{pmatrix}. \quad (45)$$

Comparing equations (44) and (45), we get

$$f_4 = \partial_4 A_4 + \vec{\nabla} \cdot \vec{A} = 0; \quad (46)$$

$$f'_4 = \partial'_4 B_4 + \vec{\nabla}' \cdot \vec{B} = 0; \quad (47)$$

$$\vec{f} = \partial_4 \vec{B} - \vec{\nabla} B_4 - \vec{\nabla} \times \vec{A}; \quad (48)$$

$$\vec{f}' = \partial_4' \vec{A} - \vec{\nabla}' A_4 - \vec{\nabla}' \times \vec{B}. \quad (49)$$

In equations (46, 47), we could get $f_4 = 0$ and $f_4' = 0$ on applying Lorentz gauge conditions on electric and magnetic four potentials. We may now discuss the different cases in the following sub sections:

(i) BRADYONIC CASE

We may describe the theory of bradyons by substituting $B_4 = A_4$, $\vec{B} = \vec{A}$ in equations (40-49) and correspondingly $\partial_4 = \partial_4'$, $\vec{\nabla} = \vec{\nabla}' = \partial$. Then we get

$$\overline{D}\emptyset = F = \begin{pmatrix} \partial_4 A_4 + \vec{\nabla} \cdot \vec{A} & -(\partial_4 \vec{A} - \vec{\nabla} A_4 - \vec{\nabla} \times \vec{A}) \\ \partial_4 \vec{A} - \vec{\nabla} A_4 - \vec{\nabla} \times \vec{A} & \partial_4 A_4 + \vec{\nabla} \cdot \vec{A} \end{pmatrix}. \quad (50)$$

Using the definition of electric and magnetic fields given by

$$\vec{E} = -\frac{\partial \vec{A}}{\partial t} - \vec{\nabla} \phi_e = -\frac{\partial \vec{A}}{\partial t} - \vec{\nabla} \phi_e; \quad (51)$$

$$\vec{\mathcal{H}} = \vec{\nabla} \times \vec{A} \quad (52)$$

and imposing Lorentz-gauge condition $\partial_4 A_4 + \vec{\nabla} \cdot \vec{A} = \frac{\partial \phi_e}{\partial t} + \vec{\nabla} \cdot \vec{A} = 0$, and using $\vec{\psi} = \vec{E} - i\vec{\mathcal{H}} (i = \sqrt{-1})$, we get

$$-\partial_4 \vec{A} + \vec{\nabla} A_4 + \vec{\nabla} \times \vec{A} = -i\vec{E} + \vec{\mathcal{H}} = -i\vec{\psi}; \quad (53)$$

$$\partial_4 \vec{A} - \vec{\nabla} A_4 - \vec{\nabla} \times \vec{A} = i\vec{E} - \vec{\mathcal{H}} = i\vec{\psi}^*. \quad (54)$$

Thus equation (50) reduces to

$$\overline{D}\emptyset = F \quad (55)$$

where

$$F = \begin{pmatrix} 0 & -\vec{F} \\ \vec{F} & 0 \end{pmatrix} = i\psi^*. \quad (56)$$

Equation (56) is the split octonion form of field tensor $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ describing the electric and magnetic fields. Off diagonal vector components of split octonions (50) are the conjugates to each other and the scalar components along principal diagonal are associated with the quaternion scalars (i.e. real quaternion). Similarly, we get

$$D\bar{\emptyset} = F^T \quad (57)$$

where

$$F^T = \begin{pmatrix} 0 & \vec{F}^* \\ -\vec{F}^* & 0 \end{pmatrix} = i\psi. \quad (58)$$

and equations (55) and (57) lead to

$$\frac{1}{2}(\bar{D}\emptyset + D\bar{\emptyset}) = \begin{pmatrix} 0 & \vec{\mathcal{H}} \\ -\vec{\mathcal{H}} & 0 \end{pmatrix} = \vec{\mathcal{H}} \quad (59)$$

$$\frac{1}{2}i(\bar{D}\emptyset - D\bar{\emptyset}) = \begin{pmatrix} 0 & \vec{E} \\ -\vec{E} & 0 \end{pmatrix} = \vec{E} \quad (60)$$

where \vec{E} and $\vec{\mathcal{H}}$ are the electric and magnetic fields given by equations (51,52). Here we have the different split octonion representation in comparison to the case quaternion electrodynamics, where $\frac{1}{2}(\bar{D}\emptyset + D\bar{\emptyset})$ represents Lorentz condition (i.e. scalar part of quaternion) and $\frac{1}{2}i(\bar{D}\emptyset - D\bar{\emptyset})$ represents the vector part (pure quaternion) of a quaternion. It is obvious because the off diagonal elements of an octonion is a quaternion while those along principal diagonal are the scalars.

(ii) TACHYONIC CASE

For the description of tachyons we substitute $A_4 = B_4$, $\vec{A} = \vec{B}$ in equations (40) to (49) and get the representation of T^4 - space. In this space (one space and three time dimensions), the scalar component is $r = |\vec{r}| = (x^2 + y^2 + z^2)^{\frac{1}{2}}$ and the time gradient vector $\vec{\nabla}' = [t_x, t_y, t_z, \partial'_4 = (-ir)]$. Hence the split octonion form of four differential operator, its conjugate and the four potential of tachyons may be written as

$$D = \begin{pmatrix} \partial'_4 & -\vec{\nabla}' \\ \vec{\nabla}' & \partial'_4 \end{pmatrix} \quad (61)$$

$$\bar{D} = \begin{pmatrix} \partial'_4 & \vec{\nabla}' \\ -\vec{\nabla}' & \partial'_4 \end{pmatrix} \quad (62)$$

$$\emptyset = \begin{pmatrix} B_4 & -\vec{B} \\ \vec{B} & B_4 \end{pmatrix}. \quad (63)$$

Thus we get

$$\overline{D}\emptyset = F' = \begin{pmatrix} \partial'_4 B_4 + \vec{\nabla} \cdot \vec{B} & -(\partial'_4 \vec{B} - \vec{\nabla}' B_4 - \vec{\nabla}' \times \vec{B}) \\ \partial'_4 \vec{B} - \vec{\nabla}' B_4 - \vec{\nabla}' \times \vec{B} & \partial'_4 B_4 + \vec{\nabla}' \cdot \vec{B} \end{pmatrix}. \quad (64)$$

Now using the following expressions for tachyonic [23, 24] electric and magnetic fields in T^4 -space as

$$\vec{E}_t = -\frac{\partial \vec{B}}{\partial r} - \vec{\nabla}' \emptyset_g = -\frac{\partial \vec{B}}{\partial r} - \vec{\nabla}' B_4; \quad (65)$$

$$\vec{H}_t = \vec{\nabla}' \times \vec{B} \quad (66)$$

and imposing Lorentz gauge condition in T^4 -space as

$$\partial'_4 B_4 + \vec{\nabla}' \cdot \vec{B} = 0,$$

we get

$$-\partial'_4 \vec{B} + \vec{\nabla}' B_4 + \vec{\nabla}' \times \vec{B} = -i\vec{E}_t + \vec{H}_t = -i\vec{\psi}_t^*; \quad (67)$$

$$\partial'_4 \vec{B} - \vec{\nabla}' B_4 - \vec{\nabla}' \times \vec{B} = i\vec{E}_t - \vec{H}_t = i\vec{\psi}_t^* \quad (68)$$

where $\vec{\psi}_t = \vec{E}_t - i\vec{H}_t$ (t denotes for tachyonic representations). Then equation (64) reduces to

$$F' = \begin{pmatrix} 0 & -\vec{F}_t \\ \vec{F}_t & 0 \end{pmatrix} = i\vec{\psi}_t^*. \quad (69)$$

Equation (64) is the split octonion form of field tensor $F'_{\mu\nu} = \partial'_\mu B'_\nu - \partial'_\nu B'_\mu$ components of which describe the electric and magnetic field in T^4 -space. Similarly, we obtain

$$D\overline{\emptyset} = F'_t \quad (70)$$

where

$$F'_t = \begin{pmatrix} 0 & \vec{F}_t^* \\ -\vec{F}_t^* & 0 \end{pmatrix} = i\vec{\psi}_t. \quad (71)$$

In the same manner, we get

$$\frac{1}{2}(\overline{D}\emptyset + D\overline{\emptyset}) = \begin{pmatrix} 0 & \vec{H}_t \\ -\vec{H}_t & 0 \end{pmatrix} = \vec{H}_t \quad (72)$$

$$\frac{1}{2}i(\overline{D}\emptyset - D\overline{\emptyset}) = \begin{pmatrix} 0 & \vec{E}_t \\ -\vec{E}_t & 0 \end{pmatrix} = \vec{E}_t \quad (73)$$

where \vec{E}_t and \vec{H}_t are the electric and magnetic fields of tachyons in T^4 -space as given by equations (65,66).

(iii) DYONIC CASE

In order to reformulate dyonic field equations in terms of split octonions and its Zorn's vector matrix realization, we replace \emptyset in equation (42) by the complex (generalized) four - potential $\{V_\mu\}$ of dyons [36] in the following form

$$V = V_4 u_0 * + V_4 u_0 + V_i u_i * + V_i u_i = \begin{pmatrix} V_4 & -\vec{V} \\ \vec{V} & V_4 \end{pmatrix} \quad (74)$$

where V_4 and \vec{V} are the temporal and spatial components of generalized four - potential of dyons. These are complex quantities and their real and imaginary components are electric and magnetic constituents [36]. Split octonion four - differential operator is now defined as,

$$D = \partial_4 u_0 * + \partial_4 u_0 + \partial_i u_i * + \partial_i u_i = \begin{pmatrix} \partial_4 & -\vec{\nabla} \\ \vec{\nabla} & \partial_4 \end{pmatrix} \quad (75)$$

Operating split octonion conjugate of four - differential operator given by equation (75) on the equation (74) and using to communication relations of split octonion units, we get

$$\overline{D}V = \begin{pmatrix} \partial_4 V_4 + \vec{\nabla} \cdot \vec{V} & -(\partial_4 \vec{V} - \vec{\nabla} V_4 - \vec{\nabla} \times \vec{V}) \\ \partial_4 \vec{V} - \vec{\nabla} V_4 - \vec{\nabla} \times \vec{V} & \partial_4 V_4 + \vec{\nabla} \cdot \vec{V} \end{pmatrix} \quad (76)$$

where

$$\partial_4 V_4 + \vec{\nabla} \cdot \vec{V} = (\partial_\mu V_\mu)(u_0 * + u_0) = 0.1 = 0 \quad (77)$$

and

$$-\partial_4 \vec{V} + \vec{\nabla} V_4 + \vec{\nabla} \times \vec{V} = -i\psi * . \quad (78)$$

Thus V is the generalized split octonion potential of dyons and $\vec{\psi}$ is the complex vector field with \vec{E} and \vec{H} as the generalized electromagnetic fields of dyons. Consequently, equation (76) leads to the following split octonion wave equation for dyonic field as

$$\overline{D}V = G \quad (79)$$

where

$$G = \begin{pmatrix} 0 & -\vec{G} \\ \vec{G} & 0 \end{pmatrix} = i\vec{\psi} * . \quad (80)$$

It is to be noted from equations (46) to (49) that f_4, \vec{f}, \vec{f}' and f'_4 are the components of electric and magnetic fields described in terms of two four-potentials in internal and external spaces. If the two spaces are completely disjoint spaces then we get only \vec{f} , in external four - space since $f_4 = 0$ due to Lorentz gauge condition while \vec{f}' and f'_4 do not occur. Equations (48) and (49) shows that \vec{f} and \vec{f}' are made up from electric and magnetic fields, which is the mixing of external and internal spaces. As such equations (48) and (49) do not describe either the usual electric and magnetic fields or the generalized fields of dyons. Rather, they have the mixed behaviour, namely tachyonic dyons.

4.2 FIELD EQUATIONS

Let us write the octonion wave equation in split representation as

$$DF = J. \quad (81)$$

Here D and F are defined by equations (40) and (45) in split octonionic form and J is associated with eight dimensional current source density in split octonion representation as follows,

$$J = \begin{pmatrix} J_4 & -\vec{J} \\ \vec{J} & J_4 \end{pmatrix}. \quad (82)$$

Now using equations (40) and (45) , we get

$$DF = \begin{pmatrix} \partial_4 f_4 - \vec{\nabla} \cdot \vec{f}' & -(\partial_4 \vec{f} + \vec{\nabla} f'_4 + \vec{\nabla}' \times f') \\ \partial_4 \vec{f}' + \vec{\nabla}' f_4 + \vec{\nabla} \times \vec{f} & \partial_4 f'_4 - \vec{\nabla}' \cdot \vec{f} \end{pmatrix}. \quad (83)$$

Comparing equations (81), (82) and (83), we get

$$J_4 = \partial_4 f_4 - \vec{\nabla} \cdot \vec{f}'; \quad (84)$$

$$J'_4 = \partial'_4 f'_4 - \vec{\nabla}' \cdot \vec{f}; \quad (85)$$

$$\vec{J} = \partial_4 \vec{f} + \vec{\nabla} f'_4 + \vec{\nabla}' \times \vec{f}'; \quad (86)$$

$$\vec{J}' = \partial'_4 \vec{f}' + \vec{\nabla}' f_4 + \vec{\nabla} \times \vec{f}. \quad (87)$$

Let us discuss the following different cases of generalized field equation.

(i) BRADYONIC CASE

For the description of bradyons, let us substitute $B_4 = A_4$, $\vec{B} = \vec{A}$, $\partial'_4 = \partial_4$, $\vec{\nabla}' = \vec{\nabla} \equiv \partial$ and $J'_4 = J_4$, $\vec{J}' = \vec{J}$. Using equations (51, 52, 56) we find that equation (83) reduces to

$$DF = \begin{pmatrix} \vec{\nabla} \cdot \vec{H} + i \vec{\nabla} \cdot \vec{E} & i \frac{\partial \vec{H}}{\partial t} + i \vec{\nabla} \times \vec{E} - \frac{\partial \vec{E}}{\partial t} + \vec{\nabla} \times \vec{H} \\ -i \frac{\partial \vec{H}}{\partial t} - i \vec{\nabla} \times \vec{E} + \frac{\partial \vec{E}}{\partial t} - \vec{\nabla} \times \vec{H} & \vec{\nabla} \cdot \vec{H} + i \vec{\nabla} \cdot \vec{E} \end{pmatrix} \quad (88)$$

Comparing it with equations (81, 82), we find that equation (88) is analogous to the split octonionic form of Maxwell's equation $F_{\mu\nu,\nu} = J_\mu$ which is the covariant form of a set of following four differential equations i.e.

$$\vec{\nabla} \cdot \vec{E} = \rho_e = -i J_4, \quad \vec{\nabla} \cdot \vec{H} = 0,$$

$$\vec{\nabla} \times \vec{H} = -\vec{J} + \frac{\partial \vec{E}}{\partial t}, \quad \vec{\nabla} \times \vec{E} = -\frac{\partial \vec{H}}{\partial t}. \quad (89)$$

Hence the octonion field equation (81) is identical to the Maxwell's field equation in compact and simple split octonion form. Equation (81) may also be written in terms of potential as

$$DF = D(\overline{D}\emptyset) = D\overline{D}\emptyset = \square\emptyset = J \quad (90)$$

which is equivalent to split octonion form of covariant field equation $\square V_\mu = J_\mu$ of classical electrodynamics. As such, we have reformulated the classical electrodynamics in terms of compact, simple and consistent representation of split octonion formulations for the case of particles traveling slower than light namely bradyons.

(ii) TACHYONIC CASE

In this case if we may substitute $A_4 = B_4$, $\vec{A} = \vec{B}$, $\partial_4 = \partial'_4$, $\vec{\nabla}' = \vec{\nabla} \equiv \partial$ and $J_4 = J'_4$, $\vec{J} = \vec{J}'$. Hence, using equations (61) and (64), we get

$$DF' = \begin{pmatrix} J'_4 & -\vec{J}' \\ \vec{J} & J_4 \end{pmatrix} = J'. \quad (91)$$

This equation is the split octonionic form of Maxwell's equation $F''_{\mu\nu,\nu} = J'_\mu$ in T^4 - space where we have used the following pairs of Maxwell's equation for tachyons [[25, 26]]

$$\vec{\nabla}' \cdot \vec{E}_t = -\rho_0, \quad \vec{\nabla}' \cdot \vec{H}_t = 0,$$

$$\vec{\nabla}' \times \vec{H}_t = -\vec{J}' + \frac{\partial \vec{E}_t}{\partial t}, \quad \vec{\nabla}' \times \vec{E}_t = -\frac{\partial \vec{H}_t}{\partial t}. \quad (92)$$

Hence equation (90) is visualized as the field equation of tachyons in T^4 - space in compact, simple and consistent split octonionic formulation. It has already been concluded [23, 24, 25] that T^4 - space for tachyons plays the same role as bradyons do in R^4 - space.

(iii) DYONIC CASE

Here, we consider $\{A_\mu\}$ and $\{B_\mu\}$ as electric and magnetic four - potentials described in external space (i.e. R^4 -space) only, D is defined by equation (75) and \emptyset is replaced by V the generalized four-potential of dyons given by equation (74) . Subsequently, on using equations (75), (76) and (79), we get

$$D(\overline{D}V) = DG = J \quad (93)$$

where

$$J = J_4 u_0 * + J_4 u_0 + J_i u_i * + J_i u_i = \begin{pmatrix} J_4 & -\vec{J} \\ \vec{J} & J_4 \end{pmatrix} \quad (94)$$

is the split octonion form of generalized four - current associated with dyons. Accordingly, we get

$$D(\overline{D}V) = (D\overline{D})V = DG = J \quad (95)$$

or equivalently

$$\square V = J \quad (96)$$

where

$$\square = (\overline{D}D) = (D\overline{D}) = \partial_4^2 + \nabla^2 = \partial_i \partial_i. \quad (97)$$

Equations (93) and (95) are the split octonion equivalents of Generalized Dirac-Maxwell's (GDM) equation of dyons. Hence we may describe split octonion wave equation as the generalized field equation of dyons in compact, simple and consistent manner. Moreover, equation (93) and (95) reproduce the field equations corresponding to the dynamics of electric charge (magnetic monopole) in the absence of magnetic (electric) charge on the dyons. Equation (95) is the thus split octonion equivalent of covariant field equations $\square V_\mu = J_\mu$ of generalized fields of dyons described earlier [26].

4.3 CURRENT EQUATION

Let us discuss the octonion wave equation as the octonion current differential equations in eight - dimension as

$$DJ = S. \quad (98)$$

Here D and J are defined earlier equations (41) and (82) in their split form and $S = \square F$ is defined as

$$S = \begin{pmatrix} S_4 & -\vec{S} \\ \vec{S}' & S'_4 \end{pmatrix}. \quad (99)$$

Now using equations (41) and (46) and equation (82) , we get

$$DJ = \begin{pmatrix} \partial_4 J_4 - \vec{\nabla} \cdot \vec{J}' & -(\partial_4 \vec{J} + \vec{\nabla} J'_4 + \vec{\nabla}' \times \vec{J}') \\ \partial_4 \vec{J}' + \vec{\nabla}' J_4 + \vec{\nabla} \times \vec{J} & \partial_4 J'_4 - \vec{\nabla}' \cdot \vec{J} \end{pmatrix}. \quad (100)$$

Now using equations (98), (99) and (100) , we get the following set of equations

$$S_4 = \partial_4 J_4 - \vec{\nabla} \cdot \vec{J}'; \quad (101)$$

$$S'_4 = \partial_4 J'_4 - \vec{\nabla}' \cdot \vec{J}; \quad (102)$$

$$\vec{S} = \partial_4 \vec{J} + \vec{\nabla} J'_4 + \vec{\nabla}' \times \vec{J}'; \quad (103)$$

$$\vec{S}' = \partial_4 \vec{J}' + \vec{\nabla}' J_4 + \vec{\nabla} \times \vec{J}; \quad (104)$$

where S is the new octonion variable parameter obtained from the operation of differential operator to current. Hence we may obtain a new kind of field equation in terms of new parameter S and accordingly we may analyze the different cases for bradyons, tachyons and dyons.

Acknowledgment- The work is supported by Uttarakhand Council of Science and Technology, Dehradun. One of us OPSN is thankful to Chinese Academy of Sciences and Third world Academy of Sciences for awarding him CAS-TWAS visiting scholar fellowship to pursue a research program in China. He is also grateful to Professor Tianjun Li for his hospitality at Institute of Theoretical Physics, Beijing, China.

References

- [1] P. Jordan, J Neuman, Von and E. P. Wigner, Ann. Math., **35** , (1934), 29.
- [2] A. Cayley, Philos. Mag. **26** , (1845), 208; R. P. Graves, "Life of Sir William Rowan Hamilton", 3 volumes, Arno Press, New York, 1975; J. C. Baez, "THE OCTONIONS" , Bull. Amer. Math. Soc., **39**, (2001), 145, math. RA/0105155; J. H. Conway, "On Quaternions and Octonions: Their Geometry, Arithmetic, and Symmetry," A K Peters Ltd, Massachusetts, 2003.
- [3] L. E. Dickson, Ann. Math., **20**, (1919), 155; W. R. Hamilton, "Elements of quaternions", Chelsea Publications Co., NY, (1969); P.G.Tait, "An elementary Treatise on Quaternions", Oxford Univ. Press (1875).
- [4] A. Pais, Phys. Rev. Lett., **7**, (1961), 291;
- [5] R. Penny, Am. J. Phys., **36**, (1968), 871; Nuovo Cimento, **B3**, (1971), 95.
- [6] K. Imaeda, H. Tachibana , M. Imaeda and S.Ohta, Il Nuovo Cimento, **B100**, (1987), 53; **B105**, (1990) 1203; Bull. Okayama Univ. Sci., **A24** , (1989) 181; K. Imaeda and H.Tachibana, Il Nuovo Cimento, ; **B104** (1989) 91; Bull. Okayama Univ. Sci., **A19**, (1984) , 93; H.Tachibana; Ph.D.Thesis, Okayama Univ. Science (1990).
- [7] G. C. Joshi, Lett Nuovo Cimento, **44**, (1985), 449; A. J. Davis and G. C. Joshi, J. Math. Phys., **27**, (1986), 3036; R. Foot and G. C. Joshi, Int. J. Theor. Phys., **28** , (1989), 1449 ; Phys. Lett., **B199**, (1987), 203.
- [8] A. Gamba, J. Math. Phys., **8**, 775 (1967); Il Nuovo Cimento, **A111** , (1998) 293; M. Gogberashvili, "Octonionic Version of Dirac Equations", hep-th/0505101; J. Phys. A: Math.Gen., **39** , (2006) 7099.
- [9] G. M. Dixon, "Division Algebras: Octonions, Quaternions, Complex Numbers, and the Algebraic Design of Physics", (Kluwer, 1994); Geoffrey Dixon, "Division Algebras, (1,9)-Space-time, Matter-antimatter Mixing" arXiv:hep-th/9303039.
- [10] S. Okubo, "Introduction to Octonion and Other Non-Associative Algebras in Physics", Montreal Memorial Lecture Series in Mathematical Physics, 2, Cambridge University Press, 1995; C.H. Barton and A. Sudbery, "MAGIC SQUARES OF LIE ALGEBRAS" arXiv:math.RA/0001083.
- [11] R. Dündarer, F. Gürsey and C. Tze, J. Math. Phys., **25**, (1980), 1496; Nucl.Phys., **B266** , (1986) , 440; F. Gürsey and C Tze, "On the Role of Division, Jordan and Related Algebras in Particle

- Physics", World Scientific, River Edge, NJ, 1996; M Günaydin and F Gürsey, Lett Nuovo Cimento, **6**,(1973), 407; M. Koca and N. Ozdes, J. Phys. A22, (1989), 1469; S. Catto, "Exceptional Projective Geometries and Internal Symmetries", hep-th/0302079; .
- [12] M. Günaydin and F. Gürsey, J. Math. Phys., **14**,(1973), 1651.
 - [13] M. Günaydin and F. Gürsey, Phys. Rev., **D9**,(1974), 3387.
 - [14] G. Domokos and S. Domokos-Kovesi, Phys. Rev., **D19**,(1979), 2984; J. Math. Phys., 18,(1978)
 - [15] . K.Morita ; Prog. Theor. Phys., **65**,(1981), 787; **73**, (1985), 999; B. A. Bernevig, J. Hu, N. Toumbas, and S.Cheng Zhang, "The Eight Dimensional Quantum Hall Effect and the Octonions", arXiv:cond-mat/0306045.
 - [16] C.A. Manogue and A. Sudbery, Phys. Rev., D40 , 4073(1989); T. Kugo and P. Townsend: Nucl. Phys., **B221**, (1983), 357; J. M. Evans, Nucl. Phys., **B298**, (1988), 92; J. Lukierski and F. Toppan, Phys. Lett., **B584** , (2004), 315 ; **B539** , (2002), 266 ; K. S. Abdel-Khalek Mostafa, "Ring Division Algebras, Self Duality and Supersymmetry", arXiv:hep-th/0002155; J. Schray, "Octonions and Supersymmetry", Ph.D. thesis, Department of Physics, Oregon State University, 1994; M. J. Duff, Ed., "The World in Eleven Dimensions: Supergravity, Super membranes and M-Theory", Institute of Physics Publishing, Bristol, 1999. MR 2001b:81002 and references cited therein.
 - [17] A. M. Buoncristiani, J. Math. Phys., **14**, (1973), 849.
 - [18] S. Marques, and C. G. Oliveita, J. Math. Phys., **26**, (1985), 3131.
 - [19] S. Marques and C G Oliveira, J. Math. Phys., **29**, (1988), 2127.
 - [20] S. Marques- Bohham, J. Math. Phys., **32**, (1991), 1383.
 - [21] S. Marques - Bonham, J. Math. Phys., **31**,(1990), 1478.
 - [22] S. Marques, C. G .Oliveira., Phys. Rev., **D36**,(1987), 1716.
 - [23] H. C. Chandola and B. S. Rajput, J. Math. Phys, **26**,(1985), 208.
 - [24] M. C. Pant, P. S. Bisht, O. P. S. Negi and B. S. Rajput, Cand. J. of Phys. **78**, (2000), 1.
 - [25] O. P. S. Negi, et al., Phys. Lett., **B105**,(1981), 281.
 - [26] O. P . S. Negi, and B. S. Rajput, Phys. Lett. **B113**,(1982), 183; Lett. Nuovo Cimento **34**,(1982), 180; Indian J. Pure and Appl. Phys., **20**,(1982), 472.
 - [27] H. C. Chandola and B. S. Rajput, Nuovo Cimento **B72**, (1982), 21; Indian J. Pure and Appl. Phys. **21**,(1983), 651.
 - [28] E. Recami and R. Mignani, Riv. Nuovo Cimento, **4** , (1974), 209; Lett. Nuovo Cimento **16**,(1976), 449.
 - [29] G. Dattoli and R. Mignani, Lett. Nuovo Cimento **22**,(1978), 65; V. Vysin, Lett Nuovo Cimento **22**, (1978), 76.
 - [30] E. Recami, Riv. Nuovo Cimento, **9** , 1-178 (1986); Found. Phys., **31** , (2001) 1119; PHYSICA, **A252** , (1998), 586.

- [31] G. D. Maccarrone and E. Recami, Lett Nuovo Cimento **34**, (1982), 251; **28**, (1980), 151 ; P. Caldirola, G. D. Maccarrone and E. Recami, Lett Nuovo Cimento, **29** , (1980), 241.
- [32] H. C. Chandola and B. S. Rajput, Indian J. Pure and Appl. Phys. **22**, (1984), 1.
- [33] H. C. Chandola and B. S. Rajput, Lett. Nuovo Cimento **40**, (1984), 277.
- [34] P. A. M. Dirac, Proc. Roy. Soc. London, **A133**, (1931), 60; Phys. Rev., **74** , (1948), 817; G. 't Hooft, Nucl. Phys., **B79**, (1974) , 276; A. M. Polyakov, JETP Lett., **20**, (1974) 194.
- [35] J. Schwinger, Science, **165** , (1978), 757; Phys. Rev., **144**, (1986), 1087; **151**, (1966), 1048; **151**, (1966), 1055; **173**, (1968); 1536; D. Zwanzinger, Phys. Rev., **176**, (1968) , 1489; Phys. Rev., **D3**, (1971), 880; B. Julia and A. Zee, Phys. Rev., **D11**, (1975), 2227; N. Craigie, "Theory and Detection of magnetic Monopoles in Gauge Theories", World Scientific, Singapur, 1986.
- [36] P. S.Bisht, O.P.S.Negi and B.S.Rajput, IL Nuovo Cimento, **A104** , (1991), 337; Prog. Theo. Phys., **85**, (1991), 157 .
- [37] Shalini Bisht, P. S. Bisht and O. P. S. Negi, Nuovo Cimento, **B113**, (1998), 1449; B.S. Rajput, S. R. Kumar and O.P.S.Negi, Lett. Nuovo **34**, (1982), 180 ; **36**, (1983), 75; V. Majernik, PHYSICA, **39**, (2000), 9; V. V. Kravchenkov, "Applied Quaternion Analysis", Helderman Verlag, Germany (2003).
- [38] Z. K. Silagadze,"Feynman's derivation of Maxwell's equations and extra dimensions", hep-th/0106235; Annalesde la Foundation Louis de Broglie, **27**, (2002) 241.
- [39] P. S. Bisht and O. P. S. Negi, Indian Journal of Pure and Applied Phys., **32** , (1994), 193; **31** , (1993), 292; Shalini Dangwal, P.S.Bisht and O.P.S.Negi, "Octonionic Gauge Formulation for Dyonic Fields", hep-th/0608061, Had. J., (2007) (In press); "Unified Split Octonion Formulation of Dyons", hep-th/0607209.